



AU9178374

(12) PATENT ABRIDGMENT (11) Document No. AU-B-78374/91
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 636762

- (54) Title
ACARICIDAL COMPOSITION
- International Patent Classification(s)
(51)^s **A01N 031/08 A01N 053/00**
- (21) Application No. : **78374/91** (22) Application Date : **14.06.91**
- (30) Priority Data
- (31) Number (32) Date (33) Country
2-158497 15.06.90 JP JAPAN
- (43) Publication Date : **19.12.91**
- (44) Publication Date of Accepted Application : **06.05.93**
- (71) Applicant(s)
SUMITOMO CHEMICAL COMPANY, LIMITED
- (72) Inventor(s)
HAYATO TESHIMA
- (74) Attorney or Agent
WATERMARK PATENT & TRADEMARK ATTORNEYS , Locked Bag 5, HAWTHORN VIC 3122
- (57) Claim

1. A method for controlling house dust mites, which comprises applying an acaricidally effective amount of 3-methyl-4-isopropylphenol and an inert carrier or diluent to a locus where house dust mites inhabit.

636702

Form 10

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952-69

COMPLETE SPECIFICATION

(ORIGINAL)

Class

Int. Class

Application Number:

Lodged:

Complete Specification Lodged:

Accepted:

Published:

Priority:

Related Art:

Name of Applicant:

SUMITOMO CHEMICAL COMPANY, LIMITED

Address of Applicant:

5-33, Kitahama 4-chome, chuo-ku, Osaka, Japan

Actual Inventor:

HAYATO TESHIMA

Address for Service:

WATERMARK PATENT & TRADEMARK ATTORNEYS.
LOCKED BAG NO. 5, HAWTHORN, VICTORIA 3122, AUSTRALIA

Complete Specification for the invention entitled:

ACARICIDAL COMPOSITION

The following statement is a full description of this invention, including the best method of performing it known to

US

ACARICIDAL COMPOSITION

The present invention relates to acaricidal compositions for controlling house dust mites.

5

Recently, with our living environment becoming comfortable owing to change in the style of buildings and spread of air conditioning systems, house dust mites breed much and not only they give unpleasant feeling, but also they cause swelling up of sting and besides, bronchial asthma and allergy. Thus, house dust mites give rise to serious social problems.

Hitherto, as acaricidal compositions for controlling house dust mites, organic phosphorous acaricides such as fenitrothion, fenthion, dichlorvos and diazinon, carbamate acaricides such as propoxur and carbaryl, and pyrethroid acaricides such as d-resmethrin and permethrin have been directly sprayed onto tatami mats or carpets in the form of aerosol, emulsion or oil solution, or impregnated into carpets or smoked, or used as antitub sheet.

However, these conventional acaricides are not satisfactory in toxicity, smell and controlling spectrum.

Under the circumstances, the inventors have conducted intensive research in an attempt to develop acaricidal compositions low in toxicity and exhibiting superior effect against various mites, As a result, it has been surprisingly found that 3-methyl-4-isopropylphenol can be used as an active ingredient of acaricidal compositions which meet the above objects and have remarkable controlling effect. The present invention has been accomplished after additional investigation thereon.

That is, the present invention relates to an acaricidal composition which contains 3-methyl-4-isopropylphenol (hereinafter referred to as "MIP") as an active ingredient.

5

MIP is prepared by converting m-cresol to an isopropyl derivative and has been used as fungicides for cosmetics or the like.

10

Mites which can controlled by the acaricidal compositions of the present invention include, for example, Dermatophagoides spp. such as Dermatophagoides farinae and Dermatophagoides pteronyssinus; Acaridae such as Tyrophagus putrescentiae and Aleuroglyphus ovatus; Glycyphagus spp. such as Glycyphagus privatus, Glycyphagus domesticus, and Glycyphagus destructor; Cheyletidae such as Cheyletus malaccensis and Cheyletus fortis; Tarsonemidae; Haplochthonius simplex; Macronyssidae such as Ornithonyssus bacoti; and mites which are parasitic on animals such as Ornithonyssus sylviarum and Dermanyssus gallinae.

20

Ordinarily, MIP is mixed with auxiliaries for preparation such as solid carriers, liquid carriers, gaseous carriers, surface active agents and film-forming agents and used in the formulations of emulsifiable concentrate, dust, granule, oil solution, aerosol, sheet and coating agent. These preparations contain 0.1 - 20 % by weight of MIP as an active ingredient.

30

Carriers used for formulation include, for example, kaolin clay, bentonite, talc and diatomaceous earth as solid carriers; aliphatic hydrocarbons such as kerosine and lamp oil, halogenated hydrocarbons such as dichloroethane, trichloroethane and carbon tetrachloride, and alcohols such as methanol, ethanol, isopropanol,

35

ethylene glycol and 2-ethoxyethanol as liquid carriers;
and gaseous carriers such as Freon [®] gas (fluorinated
hydrocarbon), LPG (liquefied petroleum gas) and dimethyl
ether. Surface active agents include, for example,
5 anionic surface active agents such as alkylsulfates and
alkyl(aryl)sulfonates and nonionic surface active agents
such as polyoxyethylene alkyl ethers and polyoxyethylene-
sorbitan fatty acid esters. Furthermore, film-forming
agents include, for example, cellulose derivatives, vinyl
10 resins, polyester resins and urethane resins.

It is possible to give polyfunctionality to
the acaricidal composition of the present invention or
enhance the effect thereof by adding thereto
15 insecticides, acaricides, synergists, pest repellents,
bactericides, fungicides, perfumes and colorants, for
example, pyrethroid compounds such as permethrin and
d-phenothrin.

20 Examples of usable insecticides, acaricides,
bactericides and fungicides are shown in the following
Table 1.

Dosage of the acaricidal composition containing
25 MIP as an active ingredient is preferably such that the
active ingredient is present in an amount of at least
100 mg per 1 m², preferably 100 mg to 10 g per 1 m², of
area in which mites should be controlled.

30

35

Table 1

Mark for compound	General name	Chemical name
(A)	Cypermethrin	α -cyano-3-phenoxybenzyl dl-cis,trans-3-(2,2-dichloro- vinyl)-2,2-dimethylcyclo- propanecarboxylate
(B)	Cyphenothrin	(RS)- α -cyano-3-phenoxybenzyl (1R)-cis,trans-chrysanthemate
(C)	Fenpropathrin	α -cyano-3-phenoxybenzyl 2,2, 3,3-tetramethylcyclopropane- carboxylate
(D)	Permethrin	3-phenoxybenzyl dl-cis,trans- 3-(2,2-dichlorovinyl)-2,2- dimethylcyclopropanecarboxyl- ate
(E)	d-Phenothrin	3-phenoxybenzyl (1R)-cis, trans-chrysanthemate
(F)	d-Resmethrin	5-benzyl-3-furylmethyl (1R)- cis,trans-chrysanthemate
(G)	Cyfluthrin	α -cyano-4-fluoro-3-phenoxy- benzyl 3-(2,2-dichlorovinyl)- 2,2-dimethylcyclopropane- carboxylate
(H)	Empenthrin	(RS)-1-ethynyl-2-methyl-2- pentenyl (1R)-cis,trans- chrysanthemate

Table 1 (continued)

Mark for compound	General name	Chemical name
(I)	Prallethrin	(S)-2-methyl-4-oxo-3-(2-propynyl)cyclopent-2-enyl (1R)-cis,trans-chrysanthemate
(J)	Ethofenprox	2-(4-ethoxyphenyl)-2-methylpropyl m-phenoxybenzyl ether
(K)	S-421	octachlorodipropyl ether
(L)	IBTA	isobornyl thiocynoacetate
(M)	Sanplas	3-bromo-2,3-diiodo-2-propenyl ethyl carbonate
(N)	TBZ	2-(4-thiazolyl)benzimidazole

When MIP is used in combination with pyrethroid compounds such as permethrin and d-phenothrin, weight ratio of MIP and pyrethroid compound is normally about 1:1 - about 30:1.

5

In that case, dosage of MIP and pyrethroid compound is 100 mg to 10 g and 100 mg to 1 g per 1 m² respectively.

10

The present invention will be explained in more detail by the following formulation examples and test examples. The present invention is not limited thereto.

15

In the following formulation examples and test examples, part is by weight and comparative compounds are shown by the marks of compounds in Table 1.

Formulation Example 1

20

An oil solution is prepared by mixing 0.2 part of MIP, 2 parts of kerosine and 97.8 parts of illuminating kerosine.

Formulation Example 2

25

An emulsifiable concentrate is prepared by mixing 10 parts of MIP, 14 parts of polyoxyethylene styrylphenyl ether, 6 parts of calcium dodecylbenzene-sulfonate and 70 parts of kerosine.

Formulation Example 3

30

A dust formulation is prepared by mixing 1 part of MIP, 89 parts of kaolin clay and 10 parts of talc.

Formulation Example 4

35

A granule formulation is prepared by well grinding and mixing 5 parts of MIP, 1 part of synthetic hydrated silicon oxide, 2 parts of calcium lignin sulfonate, 30 parts of bentonite and 62 parts of kaolin

clay, well kneading the mixture with addition of water, and then granulating and drying the kneaded product.

Formulation Example 5

5 One part of MIP, 0.3 part of compound (D) or (E), 7 parts of kerosine and 31.7 parts of deodorized coal oil are mixed and dissolved and filled in an aerosol container. After fitting a valve to the container, 60 parts of a pressure carrier (liquefied petroleum gas) is
10 pressure filled through the valve portion to obtain an aerosol.

Formulation Example 6

15 Prior to making a carpet composed of piles made of polypropylene, a first base fabric (nonwoven fabric), a packing agent comprising ethylene vinyl alcohol and a second base fabric made of jute, the piles are kneaded with MIP in an amount of 5.0 g/m^2 . Thus, a carpet
20 inhibited from breeding of house dust mites is obtained.

Formulation Example 7

25 A mixture of MIP and compound (D) or (E) is diluted with acetone and the diluted mixture is dropped on and impregnated in a nonwoven fabric at 5.0 g/m^2 of MIP or at 0.5 g/m^2 of compound (D) or (E) and then this
30 is dried to obtain a sheet inhibited from breeding of house dust mites.

Test Example 1

35 The emulsifiable concentrate obtained in Formulation Example 2 was diluted to a given concentration and uniformly coated on a filter paper (5 cm x 10 cm). After subjected to air-drying, the filter paper was folded in half and both edges were pasted. Twenty testing
40 mites (Tyrophagus putrescentiae and Dermatophagoides fariane) were introduced from the upper opening and the opening was sealed by a metallic clip. This was kept

at room temperature for 24 hours. After lapse of 24 hours, state of the mites was observed and mortality of the mites was obtained. Effects against Tyrophagus putrescentiae and Dermatophagoides farinae are shown in Table 2.

Test Example 2

Aerosol obtained in Formulation Example 5 or in accordance with Formulation Example 5 was uniformly sprayed onto a filter paper (30 cm x 30 cm) for 3 seconds and dried. Then, this filter paper was cut to 5 cm x 10 cm. This was folded in half and both edges thereof were pasted. Twenty testing mites (Tyrophagus putrescentiae and Dermatophagoides farinae) were introduced from upper opening and the opening was sealed by a metallic clip and this was kept at room temperature for 24 hours. After 24 hours, state of the mites was observed and mortality was obtained. Effects of Tyrophagus putrescentiae and Dermatophagoides farinae are shown in Table 3.

Test Example 3

A sheet inhibited from breeding of mites which was obtained in Formulation Example 7 or in accordance with Formulation Example 7 was cut to 5 cm x 5 cm and thereon was put 0.5 g of a medium adjusted to a given density (about 200 mites/g for Tyrophagus putrescentiae and about 500 mites/g for Dermatophagoides farinae). This was kept at room temperature for 2 weeks. After 2 weeks, the number of surviving mites in the medium was counted by saturated aqueous sodium chloride solution floatation method. This was compared with the number of surviving mites in control section and inhibition rate of reproduction was obtained. Effects against Tyrophagus putrescentiae and Dermatophagoides farinae are shown in Table 4.

Table 2

Dosage (g/m ²)	Mortality (%)	
	<u>Tyrophagus</u> <u>putrescentiae</u>	<u>Dermatophagoides</u> <u>farinae</u>
1.0	75	90
5.0	100	100
10.0	100	100
Untreated	0	0

Table 3

Compound	Concentration (W/W%)	Mortality (%)	
		<u>Tyrophagus</u> <u>putrescentiae</u>	<u>Dermatophagoides</u> <u>farinae</u>
MIP	1	75	83
MIP/(D)	1/0.3	100	100
MIP/(E)	1/0.3	100	100
(D)	0.3	38	40
(E)	0.3	40	45
Untreated	-	0	0

Table 4

Compound	Dosage (g/m ²)	Inhibition rate of reproduction (%)	
		<u>Tyrophagus</u> <u>putrescentiae</u>	<u>Dermatophagoides</u> <u>farinae</u>
MIP	5.0	85	92
MIP/(D)	5.0/0.5	100	100
MIP/(E)	5.0/0.5	100	100

As shown in the above examples, 3-methyl-4-isopropylphenol of the present invention exhibits excellent mite-controlling effect. Therefore, it can be spread, sprayed or coated onto tatami mat, carpet, sofa, and beddings such as mattress, bedclothes and pillow or spread, sprayed or placed in closets and warehouses.


10



15



20



25

30

35

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for controlling house dust mites, which comprises applying an acaricidally effective amount of 3-methyl-4-isopropylphenol and an inert carrier or diluent to a locus where house dust mites inhabit.
2. An acaricidal composition for controlling house dust mites which comprises an acaricidally effective amount of 3-methyl-4-isopropylphenol and a pyrethroid compound as an active ingredient, and an inert carrier or diluent.
3. The acaricidal composition according to claim 2, wherein the pyrethroid compound is permethrin.
4. The acaricidal composition according to claim 2, wherein the pyrethroid compound is d-phenothrin.
5. A method for controlling house dust mites, which comprises applying an acaricidally effective amount of 3-methyl-4-isopropylphenol to an area affected or likely to be affected by house dust mites.
6. A method as claimed in claim 5 which includes the application of a pyrethroid compound as an additional active ingredient.
7. A method as claimed in claim 6 in which the pyrethroid compound is permethrin or d-phenothrin.

DATED this 23rd day of February, 1993

SUMITOMO CHEMICAL COMPANY LIMITED

WATERMARK PATENT & TRADEMARK ATTORNEYS
THE ATRIUM 290 BURWOOD ROAD
HAWTHORN VICTORIA 3122
AUSTRALIA



ABSTRACT
ACARICIDAL COMPOSITION

An acaricidal composition for controlling house
5 dust mites which comprises an acaricidally effective
amount of 3-methyl-4-isopropylphenol, optionally containing
a pyrethroid compound.

10

15

20

25

30

35